

Floral morphometric analysis of *Prosopis affinis* Spreng. (Fabaceae) suggests flexibility of the reproductive system in isolated populations within the Brazilian Pampa

C. Carvalho de Gusmão Lôbo, V.M. Stefenon

Carvalho de Gusmão Lôbo C., Stefenon V.M., 2018. Floral morphometric analysis of *Prosopis affinis* Spreng. (Fabaceae) suggests flexibility of the reproductive system in isolated populations within the Brazilian Pampa. Ann. For. Res. 61(1): _-_.
_-.

Abstract. *Prosopis affinis* is native tree of the Brazilian Pampa, recognized as an important species to local culture and economy. Despite the ecological and the economic potential of this tree, studies in Brazil are limited to descriptions of the occurrence sites of *P. affinis*. Aiming to determine the existence of diversification at floral and/or at population levels for this species, morphological characteristics of inflorescences and flowers of *Prosopis affinis* growing in the Pampa biome were measured, while the pollen/ovule ratio and the outcrossing index were determined as an indirect estimate of the species breeding system. No clear reproductive specialization related to flower position in the inflorescence was observed, although basal flowers in the inflorescence presented reduction of the androecium. The results also showed that *P. affinis* presents a breeding system intermediary between facultative autogamy and xenogamy, with upper stamens supporting self-pollination, reduced stamens supporting outcrossing and the likely occurrence of an evolutionary shift at geographical level. These findings highlight the importance of conserving the remnants of *P. affinis* in southern Brazil and further investigate the pollination biology, the breeding system and the genetic diversity of this species, promoting its conservation and better understanding the effect of forest fragmentation in the Pampa biome. Further analyzes comparing isolated and larger continuous populations, as well as estimations of the real outcrossing rate of *P. affinis* through genetic studies and controlled experiments to evaluate auto-compatibility are recommended in order to generate more information about this issue.

Keywords Inhanduvá, outcrossing, plant reproduction, selfing

Authors. Valdir Marcos Stefenon - Universidade Federal do Pampa, Brazil; Camila Carvalho de Gusmão Lôbo - Universidade Federal do Rio Grande do Sul, Brazil.

Manuscript received February 2nd, 2018; revised March 22, 2018; accepted March 27, 2018; online first March 28, 2018.

Introduction

Adaptive implications of floral traits are related to pollination, breeding and mating system, and to the evolution of plant species (Johnson & Steiner 2000), even affecting speciation rates (Sargent 2004). These adaptive floral traits can vary spatiotemporally as a result of distinctive selective forces over differences in plant-pollinator interactions generating a complex pattern of floral adaptation between plant populations (Thompson 1999, Etcheverry et al. 2008). In this way, spatial variation in critical floral traits related to plant reproduction could lead us to infer important features of plant life history associated with reproduction process.

The worldwide-spread genus *Prosopis* L. (Fabaceae) is one of the most important plant taxa providing firewood, timber for local community, as well as edible fruits and forage for wildlife and domestic animals in arid and semi-arid regions (Burkart 1976). However, the deficiency of information about pollination, breeding and mating systems of species from this genus challenges the development of conservation and management strategies for species with economical and ecological importance.

Studies about the breeding system of seven species from section Algarobia of genus *Prosopis* (*P. glandulosa*, *P. velutina*, *P. chilensis*, *P. nigra*, *P. alba*, *P. flexuosa* and *P. ruscifolia*) revealed from total outcrossing (*P. nigra*) to auto-compatible (*P. alba*) species with up to 28% of selfing (Bessega et al. 2000).

A useful predictor of breeding system in plants is the ratio of pollen grains to ovules (P/O ratio), a methodology validated by several studies (Cruden 1977, Chouteau et al. 2006, Hentrich et al. 2010) and considered more effective than other morphological characteristic. The P/O ratio reflects the efficiency of pollination, as lower values of P/O suggest more efficient pollen transfer between conspecifics. This ratio was also shown to be direct related to the successional stage of a species (Cruden

1977). In addition, such proportions of pollen to ovules may be used estimate the outcrossing index (OCI), allowing the inference of autogamous breeding system in plants (Cruden 1977).

Prosopis affinis Spreng. is native to the Brazilian Pampa and recognized as an important tree species to local culture and economy (Marchiori & Alves 2011). Despite the ecological importance as a pioneer species and the economic potential of this tree, studies in Brazil are limited to descriptions of the occurrence sites of *P. affinis* (Alves & Marchiori 2010, 2011a, 2011b; Marchiori & Alves 2010, 2011; Marchiori et al. 2010). Therefore, the conservation and management of *P. affinis* requires informative data about species biology, genetics and reproduction.

Flowers of *P. affinis* are expected to follow the traits of the genus, with hermaphrodite cross-pollinated flowers, bees as the main pollinator, and floral specialization along the inflorescence. However, variation in these traits could occur in response to environment gradients or disturbances in abiotic or biotic conditions (Johnson & Steiner 2000, Busch 2011, Niet & Johnson 2012). Population fragmentation may decrease seed dispersal and affect the plant-pollinators interactions, disturbing fitness parameters (Aguilar et al. 2006, Stefenon et al. 2016) and the plant population genetic structure (Jacquemyn et al. 2012, Lemos et al. 2015, Nagel et al. 2015).

Since the middle of the XVIII century, the Brazilian Pampa has an economy based on monoculture and livestock, exploring the grass-dominated vegetation (Roesch et al. 2009). Such economic exploitation hinders the expansion of the forest formations and increases the genetic and evolutionary consequences of fragmentation in tree species within this biome (Lemos et al. 2014, Nagel et al. 2015, Stefenon et al. 2016). Thus, considering the fragmented distribution of the Brazilian populations of *P. affinis* within a disturbed matrix of agricultural lands and the indirect effects of habitat fragmentation in plant breeding sys-

tems, such isolated populations of this species might undergo a process of adaptation through transition from specialized to general pollination systems with mixed breeding system, giving the fact that pollinator availability might also oscillate. Aiming to test this hypothesis and to indirectly characterize the species breeding system, we examined the morphological characteristics of inflorescences and flowers of *P. affinis* from natural populations in the Brazilian Pampa and estimated the P/O ratio and the outcrossing index of this species. We intended to answer the following questions: (i) is there significant differentiation in floral morphology within the same inflorescence, reflecting some degree of diversification at floral level? (ii) Is there some significant differentiation in floral morphology among populations, reflecting some degree of diversification at geographical level? (iii) Are the estimations of P/O ratio and OCI characteristic of species with mixed breeding system?

Material and methods

Studied species

The genus *Prosopis* presents at least 40 species distributed in Asia, Africa and predominantly in America where 70% of the species can be found (Ribasky et al. 2009). *Prosopis affinis* is a tree species native to southern Brazil, Uruguay, and eastern Argentina (Pasiiecznik et al. 2001). This is a primarily outcrossing, insect pollinated tree species, currently classified as vulnerable in the Rio Grande do Sul State, Brazil (FZB 2014) due to the fragmented distribution and small size of the majority of the populations. Occurring exclusively in the Pampa biome, only six populations of *P. affinis* are documented in Brazil (Alves & Marchiori 2010, 2011a, 2011b; Marchiori & Alves 2010, 2011; Marchiori et al. 2010).

Study sites

Three natural populations of *P. affinis* occurring as isolated formations in disturbed environments near agricultural systems of rice, soybean and cattle ranching within the Brazilian Pampa were investigated. Inflorescences were collected during local summer (February) in populations Quarai (30°12'50"S, 56°32'7"W), São Vicente (29°42'13"S, 54°56'5"W) and Cacequi (29°47'54"S, 55°00'5"W). Vouchers were deposited in the Herbarium Bruno Edgar Irgang (HBEI) at Universidade Federal do Pampa, Campus São Gabriel. The vegetation of this area is characterized as steppic-savannah (Roesch et al. 2009) and climate is classified as Cfa (Humid subtropical climates) according to Köppen (Kuinchter & Buriol 2001).

Morphological differentiation among flowers

Total number of flowers per inflorescence ($n = 22$) was counted under a stereomicroscope using 400X magnification. Thirty flowers collected from eight individuals were sampled from basal, middle and apical portions of the inflorescence (Figure 1A), totalizing 90 flowers analyzed in each population. Floral morphology was characterized for individual flowers by measuring: (1) the height of the calyx, (2) the height of the corolla, (3) the height of the upper stamens, (4) the height of the lower stamens, (5) the height of the style, (6) the basal diameter of the calyx and (7) the diameter of the corolla (Figure 1B). All measures were obtained using a stereomicroscope with scales. A multivariate analysis (Principal Component Analysis, PCA) was employed to evaluate the morphometric differentiation among samples of the three populations. Since the floral differentiation may occur in distinct positions of the flower in the inflorescence, characterizing a reproductive distinction within the inflorescence, the correlation among all samples was evaluated through the nine morphometric measures in each of the three floral positions




Figure 1 (A) Inflorescence of *P. affinis*, showing the floral positions considered: basal, middle and apical. (B) Scheme of the floral traits measured. The double-headed lines represent the dimensions of each measured trait: yellow line: height of the corolla, green line: height of the style, blue line: height of the calyx, red line: diameter of the corolla, violet line: diameter of the calyx, black line: height of the lower stamens, pink line: height of the upper stamens.

(basal, middle and apical), totalizing 27 morphometric measures combined. The multivariate analysis was performed using the software PAST 3.04 (Hammer et al. 2001).

Estimations of reproductive parameters

The pollen/ovule ratio (P/O ratio) and the outcrossing index (OCI) were used to indirectly estimate the breeding system of *P. affinis*. The pollen-ovule ratio was determined by dividing the number of pollen grains per flower by the number of ovules per flower (Cruden 1977) for each floral position (basal, middle and apical) from flowers of populations Cacequi and Quarai (there was not buds in the inflorescences collected from São Vicente).

The outcrossing index was estimated according to Cruden (1977), as the sum of ascribed values for three floral features: diameter of the flower (corolla up to 1.0 mm = 0; 1-2 mm = 1; 2-6 mm = 2; >6 mm = 3), temporal separation of anthers dehiscence and stigma receptivity (homogamy and protogyny = 0; protandry

= 1), and spatial relationship of stigma and anthers (stigmas and anthers at the same level and possible contact between anthers and stigma = 0; stigma and anther spatially separated and unlikely contact = 1).

Statistical analysis

Statistical differences of means of the measured traits among populations and flower positions and of the P/O ration between populations were verified with Kruskal-Wallis hypothesis test and Dunn's test *a posteriori* ($\alpha = 0.05$) once checked for normality of samples distribution with Liliefors test and its homogeneity with Levene test, using the software Bioestat 5.0.

Results

Measurements of floral morphology

Prosopis affinis inflorescences presented in average 117.13 ± 29.50 yellowish, tubular hermaphrodite flowers (Figure 1A). Basal, middle and apical flowers have five lower and five upper stamens and a single style (Figure 1B). Interestingly, while the upper stamens may promote self-pollination, flower with style higher or with same size of the stamens were observed in 23.33% of basal flowers, 15.55% of middle flowers and 16.66% of apical flowers, a pattern that may avoid the occurrence of self-pollination.

The height of lower and upper stamens was significantly different ($p < 0.05$) across basal, middle and apical positions in the inflorescence in all populations (Figure 2A-C; Table 1). Stamens were shorter in the basal flowers for all populations. The longest ones were the middle flowers in populations Cacequi and




Figure 2 Mean values and variation in standard error of the measured floral traits of *Prosopis affinis* in populations Cacequi, Quaraí and São Vicente, in the Brazilian Pampa.

Quaraí (Figure 2A-B, Table 1), while the apical flowers were the longest in population São Vicente (Figure 2C, Table 1). The same pattern was observed for the height of the styles, with shorter structures in the basal position (Figure 2D-F, Table 1). However, the longest styles diverged in each population. In population Cacequi, the longest styles were found in the middle position of the inflorescence, while the longest styles in population São Vicente were observed in the apical position (Figure 2F, Table 1). In population Quaraí, flowers from the middle and apical positions presented styles

longer than the basal ones, but not statistically different from each other (Table 1).

The diameter of the calyx of basal flowers was smaller in all populations (Figure 2G-I, Table 1). In populations Cacequi and Quaraí, flowers from the middle and apical positions did not differ statistically, whilst the apical flowers are significantly larger in population São Vicente. Concerning the height of the calyx, the longest ones were found in the middle position of the inflorescence, although not different from apical flowers in population Cacequi (Figure 2H, Table 1). In populations

Table 1 Statistical resume of Kruskal-Wallis test and Dunn's post hoc of the measured floral traits of *Prosopis affinis* in populations Cacequi (Cac), Quarai (Qua) and São Vicente do Sul (SV), in the Brazilian Pampa. B-M: comparison between basal and middle flowers; B-A: comparison between basal and apical flowers; M-A: comparison between middle and apical flowers.

Characteristics	df	H			P			Z			P _z		
		Cac	Qua	SV	Cac	Qua	SV	Cac	Qua	SV	Cac	Qua	SV
Height of calyx	2	28.129	10.719	14.232	0.00	0.0047	0.0008	B-M: 5.282	B-M: 2.337	B-M: 3.64	<0.05	ns	<0.05
					B-A: 2.989	B-A: 0.81	B-A: 1.072	<0.05	ns	ns			
					M-A: 2.292	M-A: 3.147	M-A: 2.569	ns	<0.05	<0.05			
Height of corolla	2	44.052	32.895	17.685	0.00	0.0000	0.0001	B-M: 6.604	B-M: 5.712	B-M: 1.606	<0.05	<0.05	ns
					B-A: 2.876	B-A: 3.197	B-A: 4.146	<0.05	<0.05	<0.05			
					M-A: 3.728	M-A: 2.515	M-A: 2.54	<0.05	<0.05	<0.05			
Height of lower anthers	2	35.069	15.481	12.545	0.00	0.0040	0.0190	B-M: 5.801	B-M: 3.913	B-M: 2.206	<0.05	<0.05	ns
					B-A: 1.877	B-A: 2.297	B-A: 3.501	ns	<0.05	<0.05			
					M-A: 3.923	M-A: 1.615	M-A: 1.294	<0.05	ns	ns			
Height of upper anthers	2	45.753	20.565	15.402	0.00	0.0000	0.0050	B-M: 6.636	B-M: 4.516	B-M: 1.912	<0.05	<0.05	ns
					B-A: 2.199	B-A: 1.917	B-A: 3.921	ns	ns	<0.05			
					M-A: 4.437	M-A: 2.599	M-A: 2.008	<0.05	<0.05	<0.05			
Height of stigma	2	31.201	28.562	11.041	0.00	0.0000	0.0040	B-M: 5.448	B-M: 4.756	B-M: 1.544	<0.05	<0.05	ns
					B-A: 1.66	B-A: 4.487	B-A: 3.318	ns	<0.05	<0.05			
					M-A: 3.787	M-A: 0.269	M-A: 1.774	<0.05	ns	ns			
Diameter of calyx	2	50.461	10.63	49.680	0.00	0.0049	0.0000	B-M: 6.742	B-M: 2.636	B-M: 4.279	<0.05	<0.05	<0.05
					B-A: 5.109	B-A: 2.960	B-A: 6.906	<0.05	<0.05	<0.05			
					M-A: 1.633	M-A: 0.323	M-A: 2.626	ns	ns	<0.05			
Diameter of corolla	2	16.850	19.902	0.432	0.0002	0.0000	0.8050	B-M: 3.187	B-M: 3.558		<0.05	<0.05	
					B-A: 0.644	B-A: 0.548	ns	ns	ns				
					M-A: 3.832	M-A: 4.106	<0.05	<0.05	<0.05				

Note. Abbreviations: df: degrees of freedom, H: Kruskal-Wallis statistics, Z: post-hoc statistics; p: Kruskal-Wallis significance and pz: post-hoc significance.

Quaraí and São Vicente, basal and apical flowers did not differ in height of the calyx.

Measures of the corolla's diameter and height revealed no consistent pattern over populations (Figure 2J-L, Table 1). In population Cacequi, flowers of the middle position revealed larger diameter and height, while the diameter of basal and apical flowers did not differ and the height of basal flowers is the smallest (Figure 2J, Table 1). The diameter of basal and apical flowers did not differ in population Quaraí (Figure 2H, Table 1). In population São Vicente, no difference was observed for the diameter of corolla across the flower positions, while apical flowers revealed the largest height and basal and middle flowers did not differ in this population (Figure 2I, Table 1).

Among populations differentiation

Multivariate analysis (Figure 3A) revealed differentiation for some samples from each population, mainly for samples from São Vi-

cente and Quaraí, suggesting relatively differentiated trends among populations concerning the morphometric traits measured. The main characters related to this differentiation for population São Vicente were the height of the stamens in apical flowers [height of upper stamens in apical flowers (HUAA), height of lower stamens in apical flowers (HLAA) and total height of stamens in apical flowers (THA); Axis 2 in Figure 3A] and the height of the stamens in median and basal flowers for population Quaraí [height of upper stamens in median flowers (HUAM), total height of stamens in median flowers (THM), height of upper stamens in basal flowers (HUAB) and total height of stamens in basal flowers (THB); Axis 1 in Figure 3A].

Estimations of P/O ratio and OCI

Although the morphological traits suggest a tendency towards occurrence of outcrossing, the mean P/O ratios estimated in Cacequi (424.33±233.91 for basal flow-




Figure 3 (A) Multivariate analysis (PCA) of samples from populations Cacequi (red squares), São Vicente (black dots) and Quaraí (blue diamonds), based on 27 floral traits combined. (B) Pollen/ovule (P/O) ratio in flowers of *P. affinis* in basal flowers (Cac/B and SV/B), middle flowers (Cac/M and SV/M) and apical flowers (Cac/A and SV/A) from populations Cacequi (Cac) and São Vicente (SV), in the Brazilian Pampa. Dots represent outlier samples.

ers, 389.66 ± 95.99 for middle flowers and 362.02 ± 90.85 for apical flowers; Figure 3B) and in São Vicente (454.84 ± 295.60 for basal flowers, 421.04 ± 140.24 for middle flowers and 487.67 ± 193.113 for apical flowers, Figure 3B) did not differ statistically for flower positions and are intermediary between facultative autogamy and xenogamy (Cruden 1977), suggesting a flexible reproductive system. In addition, this value corresponds to P/O ratios observed in intermediary to highly disturbed landscapes with pioneer species, an environment where floral self-compatibility and/or self-pollination usually occurs, mainly compensating the low incidence of pollinators. Similarly, the estimated outcrossing index was $OCI = 2$ (corolla with 2-6 mm = 2; homogamy = 0; possible contact between anther and stigma = 0), characteristic of plants that are self-compatible and autogamous, with a limited degree of outcrossing (Cruden 1977). Plants with $OCI = 2$ includes native colonizers and plants growing in highly disturbed habitats (Cruden 1977). The obtained value of seed/ovule ratio was 0.503 ± 0.189 , pointing to medium taxa of ovule abortion.

Discussion

Differentiation in floral morphology suggests ongoing evolutionary shift but no pattern of specialization for pollination

The tubular yellow hermaphrodite flowers of *P. affinis* are expected to attract insects, mainly bees, as observed in Uruguay (Galera 2000) and proposed for other *Prosopis* species (Vossler 2014). Even though hermaphrodite flowers are common in Mimosoideae, they may be functionally andromonoecious as few *Prosopis* and *Inga* (Hoc et al. 1993, Barros et al. 2013) or gynomonocious as *Acacia* (Tybirk 1997), and floral differentiation within the inflorescence may be related to ongoing evolutionary shifts.

The reduced stamens observed in some

flowers of *P. affinis* in this study may represent a kind of sex allocation, precluding self-pollination in these flowers. Such female-male asymmetry may be related to flowers evolution considering that most of the modifications related to floral specialization are associated to inflorescence positions and size of gynoeceum (Diggle 2003).

In turn, morphological characteristics of calyx and corolla are direct related with pollinator attraction. While height of stamens revealed evidences of an evolutionary shift, no pattern of specialization related to pollination can be proposed based on the measurements of the calyx and corolla performed in the Pampean populations.

Long-term isolation may have caused geographic diversification in floral morphology among populations

The regional differentiation in characteristics of the inflorescences and/or flowers at population level determines a phase of the evolutionary variation in reproductive traits across the geographic range of a species (Etcheverry et al. 2008). Thus, the pattern of differentiation among populations caused by the height of stamens may represent the beginning of such a floral specialization related to inflorescence portions in the Brazilian populations of *P. affinis*. Expansion of gallery forest species in this region of the Brazilian Pampa started about 5,000 years before present (Behling et al. 2005). Considering that *P. affinis* populations were founded in the region at that time and assuming a generation interval of 15 years for the species, more than 330 generations have passed since establishment. Considering the patterns of fragmented landscape of the Brazilian Pampa concerning forest formations (Roesch et al. 2009), much likely no connection among populations existed since then and those populations experience a considerable long time of isolation and evolutionary shift. Since the evaluated floral traits have a genetic control, the geographic and reproductive isola-

tion of the studied populations of *P. affinis* for such a long period may have contributed to the differentiation observed among populations.

P/O ratio and OCI suggest a flexible reproductive system between facultative autogamy and xenogamy

The P/O ratio estimated for *P. affinis* in this study was lower than means reported by Cruden (1977) for the Mimosoideae *Caliandra palmeri* (863.3 ± 42.3), but similar to *C. calistemon* (526.5 ± 59.0), *C. kimthii* (328.7 ± 36.6) and *C. anomala* (246.1 ± 24.5). Even though such relatively low P/O ratio is a feature of Mimosaceae, the studied populations of *P. affinis* may be following such selection regarding economy in pollen production and a relative change in sex allocation through the shortening of stamens observed in some flowers.

In comparison to other xenogamous species, Mimosoideae subfamily present low values of P/O ratio (Cruden 1977). These plants present low fecundity and invest maximal energy in pollen production. However, the reproductive benefit is reasonably high when pollination is effective. As proposed by Mazer & Delesalle (1998), autogamous flowers would evolve favoring an effective P/O ratio consistent among flowers, producing as much pollen as needed to safeguard fertilization. On the other hand, outcrossing plants may experience selection towards a variation in sex allocation, conditional to temporal changes in the richness and efficiency of pollinators (Delesalle et al. 2008).

Outlook and perspectives

Flowers of *P. affinis* present upper stamens that support self-pollination, while the presence of flowers with reduced stamens supports the outcrossing. In addition, the estimations of P/O ratios and OCI are equivalent to species with a breeding system intermediary between facultative autogamy and xenogamy. Such estimations are also in accordance with the

fragmented and disturbed agricultural matrix where populations occur, because pollinators may be less abundant and less effective, prejudicing outcrossing. In addition, since the multivariate analysis revealed flowers presenting a significant divergence among populations in relation to floral morphology, an evolutionary shift at geographical level may be in course. On the other hand, reproductive differentiation within inflorescence should be refuted since the P/O ratio in each population and overall did not differ statistically for flower positions, even though a higher number of flowers with reduction of stamens' height were observed in basal flowers, in comparison to other positions. Substantial evidence on this likely evolutionary shift might be obtained from further analyzes comparing isolated and larger continuous populations. In addition, genetic studies aiming to estimate the real outcrossing rate of *P. affinis* and controlled experiments to evaluate auto-compatibility are needed in order to get more information about this issue.

As a general conclusion, this study shows that *P. affinis* presents a breeding system intermediary between facultative autogamy and xenogamy, with upper stamens supporting self-pollination, reduced stamens supporting outcrossing and the likely occurrence of an evolutionary shift at geographical level. These findings highlight the importance of conserving the remnants of *P. affinis* in southern Brazil and further investigate the pollination biology, the breeding system and the genetic diversity of this species, in order to promote its conservation and better understand the effect of forest fragmentation in the Pampa biome.

Acknowledgements

Authors knowledge the financial support provided by CNPq/Brazil (Process 442995/2014-8), FAPERGS and PROPESQ/UNIPAMPA. We thank Dr. Rubem Samuel de Ávila Jr. by the helpful discussions and comments on an

earlier version of this manuscript and MSc. Rafael Plá Matielo Lemos by field assistance.

References

- Aguilar R., Ashworth L., Galetto L., Aizen MA., 2006. Plant reproductive susceptibility to habitat fragmentation: review and synthesis through a meta analysis. *Ecological Letters* 9: 968-980.
- Alves F.S., Marchiori J.N.C., 2010. O inhanduvá (*Prosopis affinis* Spreng.) no Rio Grande do Sul. 2 – Ocorrência natural na região do Parque do Jarau, Quaraí [The inhanduvá (*Prosopis affinis* Spreng.) in Rio Grande do Sul. 2 - Natural occurrence in the region of Jarau Park, Quaraí]. *Balduínia* 25: 1-9.
- Alves F.S., Marchiori J.N.C., 2011a. O inhanduvá (*Prosopis affinis* Spreng.) no Rio Grande do Sul. 5 – Ocorrência natural na várzea do rio Santa Maria, Rosário do Sul [The inhanduvá (*Prosopis affinis* Spreng.) in Rio Grande do Sul. 5 - Natural occurrence in the floodplain of the Santa Maria river, Rosário do Sul]. *Balduínia* 27: 1-7.
- Alves F.S., Marchiori J.N.C., 2011b. O inhanduvá (*Prosopis affinis* Spreng.) no Rio Grande do Sul. 7 – Ocorrência natural na planície de inundação do arroio Itapororó, município de [The inhanduvá (*Prosopis affinis* Spreng.) in Rio Grande do Sul. 7 - Natural occurrence in the floodplain of the Itapororó stream, municipality of Alegrete]. *Balduínia* 28: 1-7.
- Barros E.C. de O., Webber A.C., Machado I.C., 2013. Limitação de polinizadores e mecanismo de autoincompatibilidade de ação tardia como causas da baixa formação de frutos em duas espécies simpátricas de Inga (Fabaceae - Mimosoideae) na Amazônia Central [Pollinators limitation and delayed action autoincompatibility mechanism as causes of low fruit formation in two sympatric species of Inga (Fabaceae - Mimosoideae) in Central Amazonia]. *Rodriguésia*. 64: 37-47
- Behling H., Pillar V.P., Bauermann S.G., 2005. Late Quaternary grassland (Campos), gallery forest, fire and climate dynamics, studied by pollen, charcoal and multivariate analysis of the São Francisco de Assis core in western Rio Grande do Sul (southern Brazil). *Review of Palaeobotany and Palynology* 133: 235-248.
- Bessegua C., Ferreyra L., Julio N., Montoya S., Saidman B., Vilardi J.C., 2000. Mating system parameters in species of genus *Prosopis* (Leguminosae). *Hereditas* 132: 19-27.
- Busch J.M., 2011. Demography, pollination, and Baker's law. *Evolution* 65: 1511-1513.
- Burkart A., 1976. A monograph of the genus *Prosopis* (Leguminosae sub-fam. Mimosoideae). *Journal of the Arnold Arboretum* 57: 219-249; 450-455.
- Chouteau M., Barabé D., Gibernau M., 2006. Pollen-ovule ratios in some Neotropical Araceae and their putative significance. *Plant Systematics and Evolution* 257: 147-157.
- Cruden R.W., 1977. Pollen-ovule ratios: a conservative indicator of breeding systems in flowering plants. *Evolution* 31: 32-46.
- Delesalle V.A., Mazer S.J., Paz H., 2008. Temporal variation in the pollen: ovule ratios of *Clarkia* (Onagraceae) taxa with contrasting mating systems: field populations. *Journal of Evolutionary Biology* 21: 310-23.
- Diggle P.K., 2003. Architectural effects on floral form and function: a review. In: Stuessy T., Horandl E., Mayer V., (eds.), *Deep morphology: toward a renaissance of morphology in plant systematics*. Koeltz: Konigstein.
- Etcheverry A.V., Alemán M.M., Fleming T.F., 2008. Flower morphology, pollination biology and mating system of the complex flower of *Vigna caracalla* (Fabaceae: Papilionoideae). *Annals of Botany* 102: 305-316
- Fundação Zoobotânica Do Rio Grande Do Sul. 2014. Taxons da flora nativa do estado Rio Grande do Sul ameaçadas de extinção. Governo do Estado do Rio Grande do Sul [Taxons of the native flora of Rio Grande do Sul state threatened with extinction]. Government of the State of Rio Grande do Sul.
- Galera F., 2000 Los algarrobos: las especies del género *Prosopis* (algarrobos) de América Latina con especial énfasis en aquellas de interés económico [The carob trees: the species of the genus *Prosopis* (carob trees) of Latin America with special emphasis on those of economic interest]. Córdoba: UNC-Secretaría de Ciencia y Tecnología, 269 p.
- Hammer Ø., Harper D.A.T., Ryan P.D., 2001. Past: Palaeontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4: 4
- Hentrich H., Kaiser R., Gottsberger G., 2010. Floral biology and reproductive isolation by floral scent in three sympatric aroid species in French Guiana. *Plant Biology* 12: 587-596.
- Hoc P.S., Agulló M.A., Palacios R., 1994. Stylar trimorphism in four functionally andromonoecious *Prosopis* species (Mimosaceae). *Plant Systematics and Evolution* 190: 143-156.
- Jacquemyn H., De Meester L., Jongejans E., Honnay O., 2012. Evolutionary changes in plant reproductive traits following habitat fragmentation and their consequences for population fitness. *Journal of Ecology* 100: 76-87.
- Johnson S.D., Steiner K.E., 2000. Generalization versus specialization in plant pollination systems. *Trends in Ecology and Evolution* 15: 140-143.
- Kuinchter A., Buriol G.A., 2001. Clima do estado do Rio Grande do Sul segundo a classificação climática de Köppen e Thornthwaite [Climate of the Rio Grande do Sul State according to the climatic classification of Köppen and Thornthwaite]. *Disciplinarum Scientia* 2: 171-182.
- Lemos R.P.M., D'Oliveira C.B., Rodrigues C.R., Roesch L.F.W., Stefenon V.M., 2014. Modeling distribution of *Schinus molle* L. in the Brazilian Pampa: insights on vegetation dynamics and conservation of the biome. *Annals of Forest Research* 57: 205-214.

- Lemos R.P.M., D'Oliveira C.B., Stefenon V.M., 2015. Genetic structure and internal gene flow in populations of *Schinus molle* (Anacardiaceae) in the Brazilian Pampa. *Tree Genetics & Genomes* 11: 75
- Marchiori J.N.C., Alves F. da S., 2010. O Inhanduvá (*Prosopis affinis* Spreng.) no Rio Grande do Sul. 1 – Embasamento fitogeográfico e pendências terminológicas [The Inhanduvá (*Prosopis affinis* Spreng.) in Rio Grande do Sul. 1 - Phylogeographic background and terminological issues]. *Balduinia* 24: 1-11.
- Marchiori J.N.C., Alves F. da S., Paz, E.A., 2010. O Inhanduvá (*Prosopis affinis* Spreng.) no Rio Grande do Sul. 3 - Parque da Cabanha do Loreto, São Vicente do Sul [The Inhanduvá (*Prosopis affinis* Spreng.) in Rio Grande do Sul. 3- Park Cabanha do Loreto, São Vicente do Sul]. *Balduinia* 25: 22-31.
- Marchiori J.N.C., Alves F. da S., 2011. O Inhanduvá (*Prosopis affinis* Spreng.) no Rio Grande do Sul. 8 – Aspectos fitogeográficos [The Inhanduvá (*Prosopis affinis* Spreng.) in Rio Grande do Sul. 8 - Phylogeographical aspects]. *Balduinia* 29: 13-20.
- Mazer S.J., Delesalle V.A., 1998. Contrasting variation within and covariation between gender-related traits in autogamous versus outcrossing species: alternative evolutionary predictions. *Evolutionary Ecology* 12: 403–425.
- Nagel J.C., Ceconi D.E., Poletto I., Stefenon V.M., 2015. Historical gene flow within and among populations of *Luehea divaricata* in the Brazilian Pampa. *Genetica* 143: 317–329.
- Niet Van der T., Johnson S.D., 2012. Phylogenetic evidence for pollinator-driven diversification of angiosperms. *Trends in Ecology and Evolution* 27: 353-361.
- Pasiecznik N.M., Felker P., Harris P.J.C., et al., 2001. The *Prosopis juliflora* – *Prosopis pallida* Complex: A Monograph. HDRA, Coventry, 172 p. UK.
- Ribasky J., Drummond M.A., Oliveira V.R., et al., 2009. Algaroba (*Prosopis juliflora*): Árvore de uso múltiplo para a região Semiárida Brasileira [Algaroba (*Prosopis juliflora*): Multiple-use tree for the Brazilian semi-arid region. Colombo. 240p.
- Roesch L.F.W., Vieira F.C.B., Pereira V.A., Schünemann A.L., Teixeira I.F., Senna A.J.T., Stefenon V.M., 2009. The Brazilian Pampa: A Fragile Biome. *Diversity* 1: 182-198.
- Sargent R.D., 2004. Floral symmetry affects speciation rates in angiosperms. *Proceedings of the Royal Society of London B* 271: 603-608.
- Stefenon V.M., Nagel J.C., Ceconi D.E., Poletto I., 2016. Evidences of genetic bottleneck and fitness decline in *Luehea divaricata* populations from southern Brazil. *Silva Fennica* 50: 1566
- Thompson J.N., 1999. Specific hypotheses on the geographic mosaic of coevolution. *The American Naturalist* 153: S1-S14.
- Tybirk K., 1997. Reproductive biology and evolution of the genus *Acacia*. *Bulletin of the International Group for the Study of Mimosoideae* 20: 45–53.
- Vossler F.G., 2014. A tight relationship between the solitary bee *Calliopsis* (*Ceroliopoeum*), *Laeta* (*Andrenidae*, *Panurginae*) and *Prosopis pollen* hosts (*Fabaceae*, *Mimosoideae*) in xeric South America woodlands. *Journal of Pollination Ecology* 14: 270-277.